

## CASE REPORT

# Minimally Invasive Beating Heart Mitral Valve Surgery

*Kalita Jyoti Prasad<sup>1</sup>, Ahangar AG<sup>2</sup>, Saikia Manuj Kumar<sup>3</sup>, Yunus Md<sup>4</sup>, Kapoor Manish<sup>5</sup>*

*Received on June 18, 2015; editorial approval on January 1<sup>st</sup>, 2016*

## ABSTRACT

*As techniques in cardiac surgery continue to improve, the minimal invasive approach is providing benefits to the cardiac surgical patient. We report a case of a 24 year old gentleman with diagnosis of rheumatic heart disease with severe mitral stenosis that underwent mitral valve replacement via minimal invasive approach on beating heart. His post operative course was uneventful and he has been on regular follow up now. The advantages and disadvantages of the procedure are discussed here.*

**Keywords:** Heart, mitral valve, thoracotomy

## INTRODUCTION

Traditional cardiac surgery generally has been performed through a median sternotomy approach. The development of technique and technology has hastened a shift towards efficient and safe minimally invasive cardiac surgery. Mitral valve surgery is one of the common operations in cardiac surgery field. As understandably, minimally invasive mitral valve surgery has advantages like decreased surgical trauma, postoperative pain, recovery time, and complications related to midline sternotomy are reduced. Several different approaches to the mitral valve has been used – namely partial lower sternotomy or right parasternal approach.<sup>1</sup> Both the approach does preserve all or part of the sternum, but it is essentially the same operation as the transsternal procedure. Yet in another approach, the surgeon can reach the heart through a right thoracotomy through the 3<sup>rd</sup> or 4<sup>th</sup> intercostal space.<sup>2</sup> Again unlike in conventional mitral valve surgery where heart is arrested with cardioplegic agent at the time of surgery, a beating heart mitral valve surgery can be performed employing certain technique and understandably with certain benefits. We describe a case in which we performed mitral valve replacement using an approach via a limited right thoracotomy on beating heart.

## THE CASE

This 24 year old gentleman felt exertional breathlessness. On evaluation he was found to have severe mitral stenosis with mitral valve area 0.7 cm<sup>2</sup>, mean mitral valve gradient of 16 mmHg with subvalvular fusion.

He was taken up for mitral valve replacement via right thoracotomy approach and on beating heart. The position of patient on table was kept supine with right thorax elevated with small pillow below the right scapula and chest. A 7-8 cm skin incision was made at mammary line

---

## Address for correspondence and reprint:

<sup>1</sup>Assistant Professor (**Corresponding Author**)

**Email:** emailtojpgk@gmail.com

**Mobile:** +918575435250

<sup>2</sup>Director; <sup>3</sup>Professor and Head, Department of Cardiothoracic and Vascular Surgery

<sup>4</sup>Additional Professor, Department of Anaesthesia

<sup>5</sup>Associate Professor, Department of Cardiology, North Eastern Indira Gandhi Regional Institute of Health and Medical Sciences, Shillong, Meghalaya.

and thoracotomy was performed via right anterior 4th intercostals space. Cardiopulmonary bypass was initiated with aortic and bicaval cannulation, and systemically cooled to 28 °C (**Figure 1**). An aortic root vent needle was inserted and connected to vent sucker at 100 mmHg. The patient was placed in the Trendelenburg position. Patient’s perfusion pressure was kept at 70 mmHg. Heart rate was maintained between 70-80 beats per minute, and right atrium was opened.




**Figure 1** Rt anterior 4<sup>th</sup> intercostal space approach & on cardiopulmonary bypass

One drop-in sucker was placed in the left atrium to maintain a bloodless operative field. Handheld retractor was used to expose the mitral valve (**Figure 2**). Anterior mitral leaflet excised, complete preservation of posterior mitral leaflet and submitral apparatus was done to maintain left ventricular geometry. A 27 ATS Medtronic prosthetic valve was seated. Rewarming was started at the time of tying the valve sutures. Prosthetic mitral valve leaflet mobility was checked. Deairing of left side of heart performed. The right atrium was closed using a standard technique.



**Figure 2** Intracardiac structures after opening of right atrium

Came out of cardiopulmonary bypass in normal sinus rhythm. Two right ventricular temporary pacing wires were put as standard protocol. Thoracotomy wound was closed in layers with intercostals drains in situ (**Figure 3**). The postoperative course was uneventful. His postoperative echo cardiogram showed normally functioning prosthetic mitral valve with normal left ventricular function. The patient discharged on 6<sup>th</sup> postoperative day.



**Figure 3** Closing the minithoracotomy surgical wound

## DISCUSSION

The first description of minimal invasive mitral valve surgery (MIMVS) did come from Navia and Cosgrove in the mid 1990s.<sup>1</sup> Since then various minimally invasive approaches for mitral valve have been reported including the parasternal, hemisternotomy, minithoracotomy and totally endoscopic approaches with a common goal of all these approaches is to avoid median sternotomy related limitations or complications.<sup>2</sup>

Beating-heart surgery is performed without stopping the heart. The circulation of blood to the heart muscle continues during the operation. Surgery on a stopped heart is common. The heart is stopped for surgery and blood to heart is reintroduced to restart the heart again. This is called reperfusion. Reperfusion can cause impairment of heart function due to ischemia-reperfusion injury. Reperfusion injury can be avoided if the heart is kept beating during surgery.

The clinical rationale for this ‘mini’ approach is to improve outcomes in valve surgery as small incisions are being used to decrease pain and trauma, improved postoperative respiratory function, reduce blood transfusion, reduced period of hospital stay and less costly, while providing the same quality of surgery. More over the patient

satisfaction is improved, since they are able to return to work and normal activity is faster.<sup>3</sup> Again MIMVS has comparable long term efficacy in measures like freedom from reoperation and long-term survival compared to standard surgery.<sup>2</sup> The most common minimally invasive approach to the mitral valve is a right minithoracotomy.<sup>4</sup> The incision is made along the 4<sup>th</sup> intercostals space. The incision extends from the parasternal border from 7 to 10 cm laterally. The mitral valve is positioned in the center of the incision although the surgical field is smaller than a median sternotomy.<sup>2</sup>

The pericardium is entered, and antegrade aortic and right atrial cannulation is performed. Patient is put on cardiopulmonary bypass and systemically cooled to 28°C to reduce metabolic demands to heart and other organs.<sup>5</sup> Keeping the ventricles empty and decompressed helps in endomyocardial coronary perfusion.<sup>5</sup> An aortic vent needle is inserted at aortic root. The left atrial approach to reach the mitral valve is common. When the left atrium is small, extension of the atriotomy over the dome of the left atrium provides an improved exposure.<sup>1</sup> When transseptal septal approach to mitral valve is chosen then both superior and inferior vena cava is cannulated separately. The inferior vena cava canula can be inserted through a different hole in lower intercostals space to increase the area of working field. This hole latter can be used for placement of intercostal drain.

The surgical field is set once patient is on cardiopulmonary bypass, then meticulous plan for final stage of operation to be carried out. When right atrial approach is chosen, it is opened after caval tapes were put down, isolating the right atrium. A few important steps to be kept in mind in this tricky and technically demanding operation. The most important stage is once the atrium is opened, it might suck air and lead to immediate massive systemic embolisation. To avoid that patient is kept in Trendelenberg position so that cerebral embolisation is minimised if at all it happens, keep the surgical field flooded with carbondioxide so that CO<sub>2</sub> embolus will get absorbed slowly, aortic root vent on optimum suction at 100 mmHg, control the patient's heart rate at around 70-80 beats/min and systemic blood pressure around 70 mmHg.<sup>4,5</sup> The left ventricular cavity should be kept filled with blood from the level of tip of mitral valve leaflets. A vent sucker put in left atrium to keep the surgical field dry through an opening in the junction of right superior pulmonary vein and left atrium. The mitral valve is examined and anterior mitral leaflet is excised.

Decalcification and release of fibrotic adhesion is done. The mitral subvalvular apparatus is preserved by reattaching it to mitral annulus. The posterior mitral leaflet is completely preserved. The preservation of mitral subvalvular apparatus is vital to maintain normal left ventricular geometry and function. A prosthetic mechanical valve is seated at mitral annulus. Rewarming is started while the surgeon ties valve sutures. Valve leaflet movement checked, left side of heart deaired and atrium is closed. Patient is repositioned, came out of cardiopulmonary bypass after putting two temporary right ventricular pacing wires for any post operative intervention if needed. The thoracic wound is closed in layers after keeping two intercostals drain, one in pericardial cavity and another in pleural cavity, temporarily.

The primary concern of minimal invasive mitral valve surgery is the incidence of neurological complications due to its possible technical limitations for adequate de-airing.<sup>5</sup> But in a systemic metaanalysis of 6 eligible studies by Seeburger et al. found no significant difference in neurological events.<sup>6</sup> A reduction in postoperative transfusion requirements is a potential advantage as significant morbidity and mortality associated with transfusions and reexploration for bleeding is minimised.<sup>5</sup> Again the comparative studies of conventional approach and minimal invasive mitral valve surgery on mortality, no study has showed a significant difference in mortality between the two approaches.<sup>2</sup> Mihaljevic et al. showed the perioperative mortality is 0.2% for the minimally invasive group and 0.3% in the sternotomy patients. Grossi et al. found 3.7% vs. 3.4% mortality between patients undergoing minimally invasive and conventional approach mitral valvesurgery.<sup>7</sup>

Who are candidates for this procedure? Patients with isolated aortic or mitral valve disease, and in some cases double valve disease, are the candidates for this procedure.<sup>8</sup> There are a few reports on isolated right coronary artery bypass grafting surgery along with this procedure.<sup>3</sup>

## CONCLUSION

Our experience in North East Indira Gandhi regional Institute of Health and Medical Sciences (NEIGRHIMS) and other eligible literature has shown the feasible alternative to the conventional full sternotomy approach to mitral valve surgery. There is a slight learning curve, but once mastered, these techniques are no different in context of amount of time needed in the procedure. There

is less perioperative morbidity; recovery is faster and cost effective. We believe that MIBHMVS is just a step in the evolution toward more minimally invasive cardiac surgical techniques that will further enhance outcomes of patients with valvular heart disease. Also this technique is excellent for atrial septal defect closure and we have been performing this operation by this technique in our institute regularly.

**Conflict of interest:** None declared.

**Consent from patient for publication:** Taken.

**Source of funding:** None declared.

**Declarations:** (1) This case article is original with the author(s) and does not infringe any copyright or violate any other right of any third parties; (2) This case article has not been published (whole or in part) elsewhere, and is not being considered for publication elsewhere in any form, except as provided herein; (3) All author(s) have contributed sufficiently in the Article to take public responsibility for it and (4) All author(s) have reviewed the final version of the above manuscript and approve it for publication.

## REFERENCE

1. Navia J Land Cosgrove D M. Minimally invasive mitral valve operations. *Annals of Thoracic Surgery* 1996;62(5):1542–1544.
2. Lucà Fabiana et al. Minimally Invasive Mitral Valve Surgery: A Systematic Review. *Minimally Invasive Surgery* Volume 03/2013;2013(1):1-10.
3. Cohn Lawrence H. Minimally invasive valve surgery. *J Card Surg* 2001;16:260–5.
4. Iribarne A, Karpenko A, Russo M J et al. Eight-year experience with minimally invasive cardiothoracic surgery. *World Journal of Surgery* 2010;34(4):611–615.
5. Michael R Petracek. Minimally Invasive Mitral Valve Surgery without Aortic Cross-Clamping. *Texas Heart Institute Journal* 2011;38(6):701–702.
6. Seeburger J, Borger MA, Falk V et al. Minimal invasive mitral valve repair for mitral regurgitation: results of 1339 consecutive patients. *European J of Cardio-Thoracic Surgery* 2008;34(4):760–765.
7. Grossi EA., LaPietra A, Ribakove GH et al. Minimally invasive versus sternotomy approaches for mitral reconstruction: comparison of intermediate-term results. *J of Thoracic and Cardiovascular Surgery* 2001;121(4):708–713.
8. Mihaljevic T, Cohn LH, Unic Det al. One thousand minimally invasive valve operations: early and late results. *Annals of Surgery* 2004;240(3):529–534.