

CASE ARTICLE

## Phthisis Bulbi Due to Ocular Trauma: A Single Case Report

**Devi Bharati<sup>1</sup>**

*Received on October 5, 2015; accepted (revised) on October 30, 2015*

### ABSTRACT

*A phthisis bulbi is a small, shrunken, non-functional eye. The affected eye may have partial vision retained in some cases; though blindness with this disorder is very common. In phthisis bulbi, the eye has a scary and shrunken appearance. Phthisis bulbi occurs as a result of trauma (commonly in case of perforating wound), accident, exposure to the radiation, tumour, eye infection or inflammation. The severity of the disorder depends upon the type and depth of injury. Scleral perforation is a very serious condition of ocular injury which leads to phthisis bulbi within a few hours. The damage to structures within the eye from any of the above mentioned causes can eventually lead to eye atrophy i.e. shrinking. In this case report a young man came with a perforating eye injury caused by a sharpened bamboo stick with lacerations around his right eye. Due to severe inflammation the ocular adnexa like sclera, cornea, conjunctiva, iris, ciliary body etc got necrosed. Eyeball became shrunken and soft with loss of vision.*

**Keywords:** *Phthisis bulbi, perforating wound, vision, blindness*

### INTRODUCTION

The eye is the organ through which an organism acquires knowledge of its surrounding environment by virtue of light reflected from or emitted by the objects within the environment. It is the photoreceptor organ which with other types of receptors allows the organism to react and understand the world around it. The human visual apparatus includes the eyeball or globe and ancillary structures such as lids and the muscles that control the eye movement. The eye is protected from mechanical injury by being enclosed in a socket, or orbit which is made up of several bones of the skull to form a four sided pyramid. Though the bony orbit provides excellent protection for the eye from blunt injury, yet the protection is not adequate. Perforating injury from glass, sharp metal fragments etc are always serious.<sup>1</sup> The injury may involve lids, lacrimal apparatus, bony orbit, the adjacent structures, the eye ball and the visual pathway. The visual loss is pronounced in penetrating injuries of the eye. The cornea bears the main brunt and lacerations of this result in iris prolapse, endophthalmitis and even phthisis bulbi. Lens changes, retinal detachment, macular and optic nerve damage are more common with blunt injury.<sup>2</sup> *Phthisis bulbi* implies, in clinical terminology, a shrunken globe, usually from ceased aqueous humor formation (phthisis meaning “wasting away”). The intraocular pressure approaches 0 mm Hg. As a consequence, the cornea becomes distorted and can develop edema and scarring, and the lens develops cataracts. Both seem to occur due to the lack of nutrition by the aqueous. In addition, edema can develop in the macula and the optic nerve head and vision suffers accordingly. Finally, cyclitic membranes and proliferative vitreal retinopathy can

---

### Address for correspondence and reprint:

<sup>1</sup>Assistant Professor  
Assam Medical College and Hospital  
Dibrugarh: 786002, Assam  
**Mobile:** 09864585459  
**Email:** devibharatidr@gmail.com

develop, resulting in total retinal detachment and scar formation. The cause of phthisis is often uveitis, either long-term or following trauma, surgery, or end-stage, heavily treated glaucoma.<sup>3</sup>

### CASE REPORT

A twenty-three year old agricultural worker from rural area came to an eye O.P.D. with a history of perforating injury to the right eye by a sharpened bamboo stick stabbed by a neighbourhood person due to some quarrel between them. He was brought by his relative to the O.P.D. 6-7 hours after the injury.



**Figure 1** Picture showing hyphaema with scleral perforation

On examination lacerated wounds were seen around his right eye. His cornea became sluggish with little perforation. Hyphaema was seen in the anterior chamber, iris became inflamed but difficult to examine due to hyphaema, scleral perforation was seen, eye lid was also lacerated, intra-ocular pressure became lower than normal but it was difficult to examine properly due to the very painful condition. Immediately after routine examination the patient was taken to the eye O.T for repair of the wounds. Then the patient was admitted in the eye ward with full treatment like injectable antibiotic, anti-inflammatory drugs, vitamins, minerals etc. He was examined regularly after the operation. The vision of his injured eye gradually came down to P.L (Perception of light) -negative. The vision of the left eye was normal-6/6. After 10 days, his eye became soft and shrunken with grayish appearance and it developed to phthisis bulbi of the right eye.

### PATHO-PHYSIOLOGY

The physiological function of the ocular adnexa as well as the intra orbital structure is always affected by trauma. The physiological alteration is determined by the severity of trauma and to restore the anatomical and physiological function of the ocular structure is the main aim of management of all cases of ocular injuries. Approximately 66% of the penetrating eye injuries are through the cornea, 10% are through sclera and the remainder through the limbal area. Small puncture or linear laceration may be self sealing and large compound laceration may be leaking with incarceration of tissues. Post traumatic uveitis may occur due to retention of foreign bodies, incarceration of uveal tissues, irritation by lens matter, haemorrhage and shallow anterior chamber. It may give rise to secondary glaucoma, plastic iridocyclitis, and shrinkage of the globe leading to phthisis bulbi and blind eye.

### DISCUSSION

Despite anatomical and physiological natural protection afforded to the eye, injuries of the eye leading to blindness of various degrees ranging from little deterioration of vision to permanent blindness are quite common.<sup>4</sup> The extent of injuries depends on the nature of trauma, extent of damage to the intraocular structures, presence of intraocular foreign bodies and the site of injury.<sup>5</sup> Grin et al categorized various types of injuries into four groups – extra ocular, ocular, intra ocular and orbital fracture. Ocular injuries made up the majority of injuries (82%) which included both perforating (34.5%) and non-perforating (50%) injuries.<sup>6</sup> Moreira et al found that most common injuries were chemical and thermal burns (24%) followed by corneal abrasion (21.9%), perforation of the globe (6.2%), subconjunctival hemorrhage (4.8%) and hyphaema (4.8%), iridodialysis accounted for 0.7%. They also found superficial corneal foreign body 3.4%, corneal delamination 1.4%, lid echymosis 3.1%, traumatic cataract 0.7%, intra ocular foreign body 0.7%. After a minimum follow up of 4 months they found that out of 38 severely injured eyes 8 (21.05%) developed traumatic cataract and 5 (13.16%) developed phthisis bulbi.<sup>7</sup> Rapoport et al found in their study that out of 1127 injured eyes, the reason for severe visual outcome were 2% due to phthisis bulbi.<sup>8</sup> Sharma et al in his study on penetrating ocular injuries found that visual acuity was severely impaired in majority of the eyes and 2% of the eyes developed phthisis bulbi.<sup>9</sup> According to Coskun M et al, after penetrating eye injuries, visual prognosis and development of phthisis

bulbi are affected significantly by the factors including anatomic localization, size of the injury, associated anterior or posterior segment pathologies, and endophthalmitis secondary to the trauma.<sup>10</sup>

## CONCLUSION

Ocular injuries may occur in all age groups but complications are more common in children and illiterate population due to lack of awareness which may lead to permanent blindness. In this case report, the patient was a young adult who sustained injury due to violence. He came for medical consultation with an ophthalmologist after 6–7 hours due to which the ocular tissues got infected leading to development of endophthalmitis and finally to phthisis bulbi. Early consultation with an eye specialist enhance the possibility of controlling the ocular morbidity. Visual impairment following ocular injury is a complex multistructural involvement which requires specialized ophthalmic training to develop the excellence for corrective and reparative ocular surgeries.

**Conflict of interest:** None

## REFERENCES

1. Albert DM, Burns WP, Scheie HG. Severe orbitocranial foreign body injury. *Am J Ophthalmol* 1965;60:1109-11.
2. Badrinath SS. Ocular trauma. *Indian J Ophthalmol* 1987;35:110-1.
3. Claes H, Dohlman MD, Donald J, D'Amico MD. Can an eye in phthisis be rehabilitated- a case of improved vision with 1-year follow-up. *Arch Ophthalmol* 1999;117(1):123-124.
4. Shukla IM, Verma RN. A clinical study of ocular injuries. *Indian J Ophthalmol* 1979 Apr;27(1):33-6.
5. Krishnan MM, Srinivasan R. Ocular injuries in union territory of Pondicherry-visual outcome. *Indian J Ophthalmol* 1988;36:86-7.
6. Grin TR, Nelson LB, Jeffers JB. Eye injuries in childhood. *Pediatrics* 1987 Jul;80(1):13-7.
7. Moreira CA, Rebeiro MD, Belfort R. Epidemiological study of eye injuries in Brazilian children. *Arch ophthalmol* 1988;106:781-84.
8. Rapoport I, Romen M, Kinek M. Eye injuries in children in Israel. *Arch Ophthalmol* 1990;108: 376–9.
9. Sharma T, Agarwal P, Gopal L, Badrinath SS, Murugesan R. Penetrating ocular trauma in children by “broomstick bows and arrows”. *Ophthalmic Surg* 1994 Mar;25(3):175-9.
10. Co'kun M, Ata° M, Akal A, Ilhan O, Keskin U, Tuzcu EA. The factors affecting the development of phthisis bulbi after penetrating eye injuries. *Ulus Travma Acil Cerrahi Derg* 2012 Jul;18(4):317-20.

## Subscription Information:

(Also available at [www.ijhrmlp.org](http://www.ijhrmlp.org))

- Founder Member of IJHRMLP will receive the journal free of cost
- Non Members and Institutions (Through Annual Subscription)
- Personal: In India, Rs. 500/Issue, Rest of the world US \$ 100/Issue
- Institutions: In India 1000/Issue, Rest of the world US \$ 200/Issue
- We accept: Bank Cheque / Demand Drafts / Online transfer