

## ORIGINAL PAPER

# Profile and knowledge of Blood Donors: A Study in a Blood Bank of a Tertiary Care Centre

*Bhattacharjee Ajanta<sup>1</sup>, Thakuria KD<sup>2</sup>, Mahanta Putul<sup>3</sup>*

*Received on .....; editorial approval .....*

## ABSTRACT

**Introduction:** *Blood is the only oxygen carrier in the body and is critical in saving human lives. The attitude, beliefs, and level of knowledge associated with blood donation will determine whether probable donors will donate blood or not. Methodology: This descriptive study with cross sectional study design will be conducted among the blood donors attending the blood bank of Tezpur Medical College and Hospital in the age group from 18 to 60 years in the department of Blood bank (Under Pathology) with due consent. Results: The response rate is 76%. The age group 20-29 years have a maximum response in terms of blood donation, i.e., 11(28.9%). Male outnumbered the female with a ratio of 2.16:1. In terms of body weight among the donors in the range of 50-59 kg have the maximum positive response, i.e., 13(34.2%). In this study, the student group comprising 15(39.5%) were the highest to respond for blood donation for the cause. Conclusion: Volunteering behavior is based on culture and determined by various factors like religion, age, sex, attitude and occupation.*

**Keywords:** *Blood Donation, Voluntary Donors, Replacement Donors, data*

## INTRODUCTION

Blood is the only oxygen transporter in the body and is crucial in saving lives. Even years of extensive research failed to find a true substitute for blood and blood components may not be available for many years.<sup>1</sup> Therefore, blood donation by humans will continue to be the major source for blood and blood components.<sup>2</sup>

A tertiary care Centre rely on the patient's relative for to donate the blood sample due to not having enough voluntary blood donations to help the needy patients.<sup>2</sup> Maximum blood donation in our hospital is on replacement basis. Blood banks pressurize the doctors, the nurses and the relatives of the patient and urge them to send replacement donors to maintain their stock. Thus the relatives of the patients are compelled

in to finding donors. Many of a time professional blood donors are brought to donate blood in a way of replacement donors. This is a very risky situation as professional donors constitute a group with high-risk behavior leading to greater chances of transfusion-transmitted diseases.<sup>3</sup>

Like in any developing country in India too, there is a dependency on family replacement and remunerated donors. Though the World Health Organization advocates that member states should establish national blood transfusion services that will operate on the basis of voluntary, non-remunerated basis<sup>4</sup>, family/replacement donors still provide more than 45% of the blood collected in our blood banks.<sup>5</sup>

This study was thus undertaken among blood donors attending the Blood bank of Tezpur Medical College and Hospital to find out their socio-demographic profile, knowledge and practice regarding blood donation.

## MATERIAL AND METHODS

This descriptive study with cross sectional study design was undertaken among the blood donors attending the blood bank of Tezpur Medical College and Hospital from December 2016 to December 2017. A total of 50 blood donors who came to donate blood during this study period in the age group from 18 to 60 years in the department of Blood bank (Under

### Address for correspondence:

<sup>1</sup>Assistant Professor

Dept. of Pathology

Email:

Mobile:

<sup>2</sup>Assistant Professor

Dept. of Physiology (Corresponding Author)

Email: kahuadas@gmail.com

Mobile: +9864051806

<sup>3</sup>Associate Professor

Dept. of Forensic Medicine

Tezpur Medical College and Hospital

This article can be cited as: Bhattacharjee Ajanta, Thakuria KD, Mahanta Putul. Profile and knowledge of Blood Donors: A Study in a Blood Bank of a Tertiary Care Centre. Int J Health Res Medico Leg Prae July-Dec 2018;4(2):85-87 DOI.....

Pathology) were randomly picked up. The final sample size thus came out to be 50.

An exit interview of the blood donors after donating blood was taken using a predesigned and pretested structured questionnaire. The study variables included age, sex, occupation and body weight. Informed consent was taken. The data thus obtained was analyzed using MS Excel.

## RESULT


A total of 50 blood donors were interviewed upon which only 38 consented to respond the questionnaire with a response rate (RR) of 76%.

**Table 1** shows the distribution of the study population according to their socio-demographic profile.


**Table 1** Distribution of the Blood Donors according to their Socio-Demographic Characteristics

Variables	Number	Percentages
<b>Age in Years</b>		
<20	2	5.3
20 - 29	11	28.9
30 - 39	10	26.3
40 - 49	0	0
50 - 59	5	13.2
≥60	10	26.3
<b>Gender</b>		
Male	26	68.4
Female	12	31.6
<b>Weight (in kg)</b>		
50 - 59	13	34.2
60 - 69	11	28.9
70 - 79	10	26.3
≥80	4	10.5
<b>Occupation</b>		
Student	15	39.5
Housewife	6	15.8
Service	4	10.5
Business	7	18.4
Farmer/Laborer	6	15.8
<b>Donor Type</b>		
Voluntary Blood Donor	19	50.0
Patient's Attendant	19	50.0

The age group of 60 and above have a maximum response, i.e., 11(28.9%) in terms of blood donation with a minimum representation from the young age group of 20 years and below, i.e., 2(5.3%). Male outnumbered the female with a ratio of 2.16:1 as shown in **Figure 1**.



**Figure 1** Gender and age variable among blood donors



**Figure 2** Variable of Body weight and occupation among blood donors

In terms of body weight among the donors in the range of 50-59 kg have the maximum positive response, i.e., 13(34.2%) and the group of 80 kg and above have the minimum (**Figure 2**). In this study, the labor class people 15(39.5%) outnumbered the other occupation for blood donation for the cause with a minimum of service holder group 4(10.5%).

## DISCUSSION

In the current study about 28.9% of the study population belonged to 20-29 years' age group. A study Uma et al<sup>6</sup> in Chennai found increased numbers of young donors who were in the age group of 18-25 years (61%). In contrast, Sampath et al<sup>7</sup> showed that 48.4% of the donors were in the age group of 26-50 years.

The number of female donors in our study was very low as compared to males with a ratio of 2.16:1. This is corroborated by similar findings in the Chennai study as well as a study by Agarwal et al<sup>8</sup>. There may be various reasons for the low

percentage of female donors like poor nutrition, low weight, and fear of pain. In a study which was done by Hollingsworth<sup>9</sup>, female donors constituted only 1% of the donor population.

In terms of body weight among the donors in the range of 50-59 kg have the maximum positive response, i.e., 13(34.2%). This positive response may be because of sound health they are enjoying within this group of body weight.

The observation that knowledge score increased with education levels ( $p<0.05$ ) seems to be logical since education would also increase awareness about all possible information related to blood donation. There is also a higher probability of having been exposed to a voluntary blood donation camp in one's educational institute since these camps are common in educational institutes.

The involvement of student group in blood donation with highest 39.5% among all group in this study is in agreement with study<sup>10</sup> carried out in Tanzania where voluntary blood donation correlated with secondary school attendance.

## CONCLUSION

The demand for blood products is ever lasting and increasing. Concurrently, blood donor enrolment becomes more and more difficult and scanty. Therefore, in this occasion volunteer blood donation should be encouraged, especially the young generation, as they can be a healthy and continuous donor. Volunteering behavior is based on culture and demographic variables and determined by various factors like age, sex, body weight and occupation, etc. Volunteer donors feel it as a responsibility to help others, regardless of personal connection. Our study found that all the respondents in the category of cultivators were willing to donate blood again if required in future. This positive feeling should be encouraged with proper knowledge regarding blood safety to meet the demand of blood requirement.

**Conflict of interest:** No conflict of interest associated with this work.

**Ethical issues:** Ethical clearance was sought from the institutional ethics committee.

**Source of funding:** None.

**Author's contribution:** We declare that this work was done

by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors.

## REFERENCES

1. Lowe KC, Ferguson E. Benefit and risk perceptions in transfusion medicine: blood and blood substitutes. *Journal of Internal Medicine* 2003;253:498–507.
2. Dutta S, Sinha D, Chatterjee S, Basu M, Misra RN. Profile and Awareness of Blood Donors: A Study In Blood Bank of a Tertiary Care Hospital of Kolkata. *Int J Pharm Sci Res* 2016;7(9):3881-86.
3. N, Thankappan KR, Kartha CC, Pal R. Analyzing Sociodemographic factors amongst blood donors. *J Emerg Trauma Shock* 2010;3:21–25.
4. The Melbourne declaration on 100% voluntary non-remunerate donation of blood and blood components,” Composed at World Blood Donor Day. Melbourne, Australia, 2009.
5. National AIDS Control Organization: Voluntary Blood Donation Program –An operational guideline; Ministry of Health and family Welfare 2007:9.
6. Uma S, Arun R, Arumugam P. The Knowledge, Attitude and Practice towards Blood Donation among Voluntary Blood Donors in Chennai. *Indian Journal of Clinical and Diagnostic Research* 2013;7:1043-1046.
7. Sampath S, Ramsaran V, Parasram S, Mohammed S, Latchman S, Khunja R, et al. Attitude towards blood donation in Trinidad and Tobago. *Transfusion Medicine* 2007;17:83-87.
8. Agarwal A, Tiwari A K, Ahuja A, Kalra R. Knowledge, attitude and practices of people towards voluntary blood donation in Uttarakhand. *Asian J Transfus Sci* 2013 Jan-Jun;7(1):59–62.
9. Hollingsworth B, Wildman J. What population influences the decision to donate blood? *Transfusion medicine* 2004;14:9-12.
10. Jacobs B, Berege ZA. Attitudes and beliefs about blood donation among adults in Mwanza Region, Tanzania. *East Afr Med J* 1995;72:345–8.